## AMENDMENT TO THE CLAIMS

Claims 1-11 (Cancelled)

Claim 12 (Currently amended). A process for the preparation of amidic cross-linked polysaccharides containing carboxy groups in which the polysaccharide is selected from the group consisting of hyaluronic acids, cabexymethyldextran carboxymethyldextran, carboxymethylcellulose, carboxymethylstarch, alginic acids, cellulose acid, N-carboxy-methyl or butyl glucans or chitosans, heparins with different molecular weights, optionally desulphated desulphated and succinylated, dermatan sulphates, chondroitin sulphates and heparan sulphates comprising (a) activating the carboxy groups of the polysaccharide in an anhydrous aprotic solvent using a suitable carboxy activating agent; (b) reacting the carboxy activated polysaccharide with a polyamine selected from the group having the formula R1-NH-A-NH-R2 wherein R1 and R2, which may be the same or different, are hydrogen, C1-C6 alkyl, phenyl or benzyl groups; A is a C2-C10 alkylene chain which may be substituted by hydroxy, carboxy, halogen, alkexy or amino groups; a polyoxyalkylene chain of the formula  $[(CH_2)_n-O-CH_2)_n]_m$ wherein n is 2 or 3 and m is an interger from 2 to 10; a C5-C7 cycloalkyl group or an aryl or heteroaryl group; and (c) recovering the resultant cross-linked polysaccharide.

Claim 13 (Previously added). A process according to Claim 12 in which the carboxy-containing polysaccharide is a hyaluronic

acid salified with a lipophilic cation; the solvent is selected from tetrahydrofuran, dimethylformamide or dimethyl sulfoxide; the carboxy activating agent is chloromethylpyridylium iodide and the polyamine is one in which A of the formula  $R_1NH-A-R_2$  is a  $C_2-C_6$  linear alkylene chain.

Claim 14 (Previously added). A process according to Claim 13 in which the polyamine, diluted in a like solvent as used in the activation step, is added to the solution of activated polysaccharide to effect the cross-linking reaction in 1-12 hours.

Claim 15 (Previously added). A process according to Claim 13 in which the recovered cross-linked polysaccharide is sulphated by reaction with a pyridine-sulfur trioxide complex.

Claim 16 (Previously added). A process according to Claim 13 in which the recovered cross-linked polysaccharide is complexed with a metal ion selected from zinc, copper and iron.

Claim 17 (Currently amended) A cross-linked polysaccharide prepared according to the process of Claim 13 characterized by various properties and physical shapes suitable for use in varying medical and veterinarian applications.